

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

Dr.Z.H.Aliyev

Institute of Soil Science and Agrochemistry of NAS of Azerbaijan, Az 1073. Baku.str. M.Raqima 5.


*Corresponding Author: Dr. Z.H.Aliyev Institute of Soil Science and Agrochemistry of NAS of Azerbaijan, Az1073. Baku.str Email: zakirakademik@mail.ru

ABSTRACT

The article examines the issues of studying the degree of susceptibility of sloping lands in Azerbaijan in the example of specific administrative-territorial units, flat areas with a slope of up to 6% (about 3.50) are concentrated in Akhsu district - 71%, and in Shemakha - almost 49%. The steepest slopes are observed in the Ismaili region, where almost 26% of the territory has a slope of 10-18%, 30% of its area slope is 18%.

Keywords: sloping slopes, arable lands, a layer, a database, a gradient class, soil, forests, pastures, geese, cartographic materials

INTRODUCTION



Spatial analysis includes operations performed on geographic data using available methods and techniques in GIS software, with a view to describing the relationships between elements of the geographical environment. The analysis can be carried out on the data, both in the vector and raster systems, and touch the geometry and attributes of the vector data.

SEARCH FOR INFORMATION IN THE DATABASE

The main operation that can be performed based on the GIS database is information retrieval. This database has a relational nature and therefore, the object designation provokes the selection of the corresponding records from the corresponding attribute table, and vice versa. The first thematic layer in the database was the Digital Elevation Model in raster format. On its basis, as a result of the transformations and calculations, information on the height of the

nos. and the slope of the terrain on the territory of Azerbaijan selected for analysis, which includes three regions: Shemakha, Akhsu and Ismayil.

VERTICAL POSITION FACTOR

The major part of the study area is in the following ranges of heights: 1) from 0 to 300 mas-23.16%, from 300 to 600 m-19.84% and 3) 600-900 m-23, 75% of the total area. Above 900 m and up to 3400 m above sea level, about 30% of its area is located (Table 1).

Of the administrative units belonging to the study area, the most highly allocated land Ismaili region. In this area 90: it is at an altitude of more than 300 meters above sea level, and sometimes the altitude is even higher than 3400 m. In Akhsu region about 73% of the earth is at an altitude up to the sea level. Shemakha district is an intermediate region, its largest area is at an altitude of 900 m above sea level-about 75%. In this area there is also a territory located below sea level-31244.66 hectares.

CLASSES OF SLOPE

In the analyzed territory, the largest area is occupied by areas with a slope of up to 6-46%. Significant areas are also in the intervals from 6 to 10% -18% and from 18% to almost 23% of the total area (Table2). Against the backdrop of

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

administrative-territorial units, gently sloping areas with a slope of up to 6% (about 3.50) are concentrated in the Akhsu district - 71%, and in Shemakha - almost 49%. The steepest slopes are

observed in the Ismaili region, where almost 26% of the territory has a slope of 10-18%, 30% of its area slope is 18%.

Table 1. The distribution of the study area (ha and%) over the altitude intervals

Height, m. h.y.m.	Regions			Togethe	
	Shemakha	Akhsuinsky	Ismailinskiy		
<0	ha	31 245	0	0	31245
	%	8,12	0,00	0,00	5
0-300	ha	69 737	73029	17334	160101
	%	18,13	72,99	8,39	23
300-600	ha	84924	11744	40470	137138
	%	22,08	11,74	19,58	20
600-900	ha	101 678	13550	48965	164193
	%	26,44	13,54	23,69	24
900-1200	ha	46975	1562	15671	64208
	%	12,21	1,56	7,58	9
1200-1500	ha	23469	162	20529	44159
	%	6,10	0,16	9,93	6
1500-1800	ha	15775	0	22305	38080
	%	4,10	0,00	10,79	6
1800-2100	ha	7885	0	22460	30345
	%	2,05	0,00	10,87	4
2100-2400	ha	2279	0	10223	12502
	%	0,59	0,00	4,95	2
2400-2700	ha	572	0	4197	4769
	%	0,15	0,00	2,03	1
2700-3000	ha	42	0	2915	2967
	%	0,01	0,00	1,41	0
3000-3300	ha	0	0	1400	1400
	%	0,00	0,00	0,68	0
>3400	ha	0	0	191	191
	%	0,00	0,00	0,09	0
Together	ha	384 582	100 047	206660	691289

CLASSES OF THE SLOPE OF THE TERRAIN, DEPENDING ON THE ALTITUDE INTERVALS

Tables (3-5) contain data that allow analyzing the distribution of the slope of the terrain along altitude intervals. This distribution was prepared for all administrative units of the area under consideration.

In the Shemakha region, a clear dependence is shown, an increase in the slope, along with an increase in altitude above sea level. Up to a height of 1200 m there are lands with a slight slope-up to 6% (3.50). They occupy 49% of the area. Territories here with a slope of 6-10% and 10-18% are located at an altitude of 400-1700

m. M.u.m.-about 23% of the area's area. Areas with a slope of more than 18% make up more than 5% of the total area and are located in the highlands. In the Aksuinsky area, the terrain with an insignificant slope of up to 6% (3.50) prevails and they are located mainly at an altitude of 300 m above sea level - this is 71% of the total area.

Areas with a slope of more than 6% are located at altitudes from 300 to 1400 m above sea level - about 29%. Clearly marked (17% of the total area) of land with a slope of 10-18% (6-100) (Table 4).

Table2. The distribution of the study area (ha and%) over the slopes of the terrain

Height, m. h.y.m	Regions				Together
		Shemakha	Akhsuinsky	Ismailinskiy	
0-6	ha	187157	71131	61504	319792
	%	48,67	71,10	29,76	46,26

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

6-10	ha	88830	7393	28770	124993
	%	23,10	7,39	13,92	18,08
10-18	ha	87641	17033	53425	158099
	%	22,79	17,02	25,85	22,87
18-27	ha	17782	4326	34060	56168
	%	4,62	4,32	16,48	8,13
27-35	ha	2855	164	20176	23195
	%	0,74	0,16	9,76	3,36
>35	ha	317	0	8724	9041
	%	0,08	0,00	4,22	1,31
Together		384582	100 047	206660	691289

In the Ismayilli region, most of the territory with a slope of <6 to 10% is located at an altitude of 300 to 900 m above sea level. Areas with a slope of 10 to 37% occur at an altitude of 3000 meters above sea level. Locations with a slope of more than 37% are located at the highest altitude, but make up only 4% of the total area of the area.

The second, from the thematic layers created in the database, is the land use of the study area. Due to the availability of cartographic materials, it was considered exclusively for the territory of the Akhsu district.

Analysis of the structure of land use in the Akhsu district showed that it is purely

Table 3. The distribution of the territory of the Shemakha district according to the grades of the terrain according to the altitude intervals

Height, m above sea level 1	slope classes%												sum	
	0-6		6-10		10-18		18-27		27-37		37-52			
	Ha	%	ha	%	ha	%	ha	%	ha	%	ha	%		
1	2	3	4	5	6	7	8	9	10	11	12	13	14	
<0	31245	8.12	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	31245	
0-100	37036	9,63	714	0,19	251	0,07	27	0,01	2	0,00	0	0,00	38030	
100-200	13223	3,44	2852	0,74	1957	0,51	372	0,10	19	0,00	0	0,00	18422	
200-300	4661	1,21	4076	1,06	3730	0,97	752	0,20	67	0,02	0	0,00	13285	
300-400	7157	1,86	5037	1,31	5548	1,44	1159	0,30	73	0,02	0	0,00	18978	
400-500	15741	4,09	7078	1,84	6609	1,72	1302	0,34	70	0,02	0	0,00	30800	
500-600	16189	4,21	8845	2,30	8644	2,25	1420	0,37	52	0,01	0	0,00	35150	
600-700	17033	4,43	11665	3,03	9814	2,55	1245	0,32	41	0,01	1	0,00	39799	
700-800	16708	4,34	12164	3,16	8375	2,18	890	0,23	24	0,01	1	0,00	38162	
800-900	7497	1,95	8050	2,09	7152	1,86	962	0,25	55	0,01	1	0,00	23717	
900-1000	4810	1,25	6280	1,63	5615	1,46	786	0,20	100	0,03	5	0,00	17596	
1000-1100	4415	1,15	4804	1,25	5480	1,43	658	0,17	97	0,03	11	0,00	15466	
1100-1200	4078	1,06	4265	1,11	4780	1,24	637	0,17	128	0,03	25	0,01	13914	
1200-1300	3402	0,88	3522	0,92	3348	0,84	424	0,11	101	0,03	18	0,00	10816	
1300-1400	1625	0,42	2665	0,69	2605	0,68	287	0,07	79	0,02	25	0,01	7286	
1400-1500	722	0,19	1880	0,49	2342	0,61	342	0,09	73	0,02	9	0,00	5368	
1500-1600	472	0,12	1765	0,46	2806	0,73	634	0,16	63	0,02	1	0,00	5741	
1600-1700	497	0,13	1161	0,30	2885	0,75	833	0,22	58	0,02	2	0,00	5437	
1700-1800	290	0,08	988	0,26	2183	0,57	1063	0,28	73	0,02	1	0,00	4597	
1800-1900	230	0,06	650	0,17	1654	0,43	1040	0,27	133	0,03	1	0,00	3708	
1900-2000	88	0,02	242	0,06	991	0,26	998	0,26	203	0,05	1	0,00	2523	
2000-2100	32	0,01	69	0,02	439	0,11	741	0,19	357	0,09	17	0,00	1655	
2100-2200	1	0,00	17	0,00	194	0,05	521	0,14	346	0,09	45	0,01	1124	
2200-2300	2	0,00	16	0,00	102	0,03	353	0,09	264	0,07	46	0,01	782	
2300-2400	5	0,00	4	0,00	28	0,01	132	0,03	156	0,04	49	0,01	373	
1	20	3	4	5	6	7	8	9	10	11	12	13	14	

agricultural. The area is dominated by arable land, which makes up about 50% and tilled areas, which include pastures, meadows and degrees - more than 23% of the total area. Agricultural lands are supplemented with vegetable gardens, orchards and vineyards, occupying 3% of the area. There are few forests and shrubs, in general, about 12%.

There are also areas completely devoid of vegetation in the Akhsu district, about 4% of them. A small part is occupied by water reservoirs, wetlands and urban and rural areas, amounting to 6.9% of the total area.

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

2400-2500	0	0,00	11	0,00	42	0,01	92	0,02	111	0,03	27	0,01	282
2500-2600	0	0,00	7	0,00	27	0,01	75	0,02	62	0,02	24	0,01	197
2600-2700	0	0,00	3	0,00	20	0,01	25	0,01	41	0,01	5	0,00	94
2700-2800	0	0,00	1	0,00	14	0,00	12	0,00	7	0,00	2	0,00	35
2800-2900	0	0,00	1	0,00	3	0,00	2	0,00	0	0,00	0	0,00	7
sum	187 157	48,67	88830	23,10	87641	22,79	17782	4,62	2855	0,74	317	0,08	384582

Table 4. Distribution of the territory of the Akhsu region according to the grade of the terrain depending on the altitude intervals

Height, m. above sea level	Slope classes%										Sum	
	0-6		6-10		10-18		18-27		27-35			
	Га	%	га	%	га	%	га	%	га	%		
<100	52626	52,60	16	0,02	13	0,01	0	0,00	0	0,00	52656	
100-200	12380	12,37	817	0,82	485	0,49	74	0,07	0	0,00	13757	
200-300	3315	3,31	1150	1,15	1830	1,83	317	0,32	5	0,01	6617	
300-400	418	0,42	693	0,69	2051	2,05	507	0,51	19	0,02	3687	
400-500	215	0,21	390	0,39	2220	2,22	834	0,83	13	0,01	3672	
500-600	273	0,27	538	0,54	2594	2,59	974	0,97	5	0,01	4384	
600-700	327	0,33	575	0,76	3338	3,34	846	0,88	20	0,02	5318	
700-800	690	0,69	1516	1,52	2689	2,69	406	0,41	22	0,02	5324	
800-900	681	0,68	988	0,99	1060	1,06	166	0,17	13	0,01	2909	
900-1000	154	0,15	254	0,28	241	0,24	79	0,08	16	0,02	775	
1000-1100	29	0,03	198	0,20	281	0,28	27	0,03	16	0,02	551	
1100-1200	23	0,02	36	0,04	127	0,13	32	0,03	17	0,02	236	
1200-1300	0	0,00	7	0,01	82	0,08	29	0,03	13	0,01	131	
1300-1400	0	0,00	4	0,00	20	0,02	5	0,00	3	0,00	31	
Sum	71131	71,1	7393	7,39	17033	17,02	4326	4,32	164	0,16	100 047	

Table 5. The distribution of the territory of the Ismayilli region according to the grades of the terrain according to the altitude intervals

Height, m. above sea level	Slope classes%										Sum		
	<6		6-10		10-18		18-27		27-37				
	Га	%	га	%	га	%	га	%	га	%			
1	2	3	4	5	6	7	8	9	10	11	12	13	14
<100	1849	0,89	321	0,16	92	0,04	0	0,00	0	0,00	0	0,00	2262
100-200	624	0,30	376	0,18	484	0,23	51	0,02	10	0,01	0	0,00	1546
200-300	10302	4,98	1937	0,94	1065	0,52	194	0,09	29	0,01	0	0,00	13,527
300-400	3303	1,60	2322	1,12	2564	1,24	288	0,14	26	0,01	3	0,00	8507
400-500	4112	1,99	2019	0,98	3240	1,57	569	0,29	30	0,01	2	0,00	9999
500-600	13759	6,66	28668	1,39	4220	2,04	1090	0,53	26	0,01	1	0,00	21964
600-700	13807	6,68	4012	1,94	4640	2,25	982	0,48	32	0,02	3	0,00	23476
700-800	8587	4,15	4068	1,97	3569	1,73	841	0,41	82	0,04	7	0,00	17154
800-900	2114	1,02	2386	1,15	2679	1,30	985	0,48	147	0,07	25	0,01	8336
900-1000	872	0,42	1190	0,58	2029	0,98	1130	0,55	349	0,17	48	0,02	5617
1000-1100	414	0,20	740	0,36	1699	0,82	1446	0,70	492	0,24	96	0,05	4887
1100-1200	194	0,09	552	0,27	1713	0,83	1776	0,86	776	0,38	158	0,08	5167
1200-1300	212	0,10	621	0,30	2015	0,98	2117	1,02	1025	0,50	191	0,09	6181
1300-1400	182	0,09	720	0,35	2553	1,24	2269	1,10	1154	0,56	237	0,11	7114
1400-1500	177	0,09	695	0,32	2440	1,18	2325	1,13	1359	0,66	276	0,13	7233
1500-1600	119	0,6	547	0,26	2499	1,21	2275	1,10	1429	0,69	376	0,18	7244
1600-1700	152	0,7	558	0,27	2440	1,18	2394	1,16	1527	0,74	455	0,22	7527
1700-1800	154	0,07	520	0,25	2490	1,21	2339	1,12	1561	0,76	448	0,24	7534
1800-1900	153	0,07	610	0,30	2743	1,32	2440	1,18	1522	0,74	533	0,26	7991
1900-2000	174	0,08	638	0,31	2682	1,30	2112	1,18	1268	0,61	604	0,29	7806
2000-2100	86	0,04	405	0,20	2355	1,14	1439	1,02	1130	0,55	573	0,28	6662
2100-2200	81	0,04	359	0,17	1723	0,83	661	0,70	1057	0,51	572	0,28	5231
2200-2300	59	0,03	246	0,12	769	0,37	2439	0,32	794	0,38	619	0,30	3148
2300-2400	7	0,00	30	0,01	156	0,08	373	0,18	682	0,33	596	0,29	1844
2400-2500	2	0,00	9	0,00	53	0,03	217	0,10	674	0,33	569	0,28	1523
1	20	3	4	5	6	7	8	9	10	11	12	13	14
2500-2600	0	0,00	2	0,00	52	0,02	232	0,11	603	0,29	562	0,27	1451
2600-2700	2	0,00	6	0,00	42	0,02	172	0,08	557	0,27	446	0,22	1223
2700-2800	1	0,00	5	0,00	65	0,03	168	0,08	509	0,25	394	0,19	1141
2800-2900	0	0,00	10	0,00	66	0,03	141	0,07	382	0,18	317	0,15	916

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

2900-3000	6	0,00	7	0,00	77	0,04	173	0,08	329	0,16	267	0,13	858
3000-3100	1	0,00	21	0,00	84	0,04	183	0,09	292	0,14	174	0,08	754
3100-3200	2	0,00	2	0,00	59	0,03	110	0,05	177	0,09	98	0,05	448
3200-3300	0	0,00	3	0,00	34	0,02	59	0,03	76	0,04	26	0,01	198
3300-3400	0	0,00	3	0,00	35	0,02	38	0,02	42	0,02	4	0,00	122
3400-3500	0	0,00	0	0,00	3	0,00	16	0,01	21	0,01	2	0,00	42
3500-3600	0	0,00	2	0,00	4	0,00	11	0,01	11	0,01	0	0,00	28
Sum	61504	29,76	28,770	13,92	53425	25,85	34060	16,48	20176	9,76	8724	4,22	206660

Table 6. Land use in Ahsuinsky district

Land use	ha	%
The Cities	1237	1,24
Pshchenitsa	49979	49,96
Gardens	1123	1,12
Vineyards	528	0,53
Arable land	52868	52,84
Pastures	12394	12,39
Green grounds	6511	6,51
Power	4708	4,71
Dry dried valleys	673	0,67
Coniferous forests	535	0,53
Deciduous forests	8451	8,45
Mixed forests	216	0,22
Marshes	844	0,84
Reservoirs	27	0,03
Stavy	25	0,02
Piece of lake	80	0,08
Bays	1137	1,14
Shrubs	2668	2,67
Soil without vegetation	248	0,25
Rocks	3827	3,38
Rural low construction	527	0,53
City low building	3919	3,92
High Urban Construction	147	0,15
Rockstones, mines	241	0,24
Together	100 047	100 00

Tables 7 and 8 provide information on the distribution of land use patterns over altitudinal intervals and grades of terrain slopes. Most agricultural land is located at a height of up to 400 m above sea level. Only pastures are located up to an altitude of 1400 meters above sea level. forest territories are in all altitude intervals. The

remaining forms of land use are located at lower altitudes (table7.).

Considering the slope of the slopes, most of the territories are in the gradient class 0-6% - almost 71%. Only pastures, green lands, as well as deciduous forests, are located on slopes with a higher slope (Table 8.).

Table 7. Land use by altitude intervals

Land use	Height, m. above sea level																
	<100		100-200		200-300		300-400		400-500		500-600		600-700		700-800		
	ha	%	ha	%	ha	%	ha	%	ha	%	ha	%	ha	%	ha	%	
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	
The Cities	231	0,23	953	0,95	53	0,05	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0
Pshchenitsa	42825	42,81	5607	5,60	690	0,69	235	0,23	0	0,00	0	0,00	7	0,01	142	0,14	
Gardens	339	0,34	701	0,70	42	0,04	0	0,00	0	0,00	0	0,00	1	0,00	41	0,04	
Vineyards	19	0,02	113	0,11	306	0,31	40	0,04	11	0,01	0	0,00	0	0,00	35	0,04	
Pastures	438	0,44	1635	1,63	461	0,46	511	0,51	691	0,69	1392	1,39	2385	2,38	2623	2,62	
Green grounds	42	0,04	596	0,60	1345	1,34	1418	1,42	1239	1,24	873	0,87	495	0,49	31	0,31	
Power	3188	3,19	1459	1,46	61	0,06	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	
DryDriedLoose Valleys	223	0,22	242	0,24	201	0,20	6	0,01	0	0,00	0	0,00	0	0,00	0	0,00	
Coniferous forests	535	0,53	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

Deciduous forests	0	0,00	93	0,09	62	0,06	421	0,42	1018	1,02	1546	1,55	1928	1,93	1901	1,90
Mixed forests	33	0,03	76	,08	104	0,10	0	0,00	0	0,00	3	0,00	0	0,00	0	0,00
Marshes	844	0,84	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00
Reservoirs	0	0,00	27	0,03	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00
Stavy	23	0,02	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	2	0,00
Piece of lake	0	0,00	80	0,08	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00
Bays	0	0,00	82	0,08	289	0,29	165	0,16	96	0,10	196	0,20	230	0,23	78	0,08
Shrubs	1110	1,11	312	0,31	1244	1,24	2	0,00	0	0,00	0	0,00	0	0,00	0	0,00
The soil layer without vegetation	138	0,14	56	0,06	53	0,05	1	0,00	0	0,00	0	0,00	0	0,00	0	0,00
Rocks	1	0,00	414	0,41	1114	1,11	904	0,90	627	0,63	372	0,37	273	0,24	136	0,14
Rural low construction	0	0,00	219	0,22	45	0,04	8	0,01	0	0,00	0	0,00	16	0,02	38	0,04
Citylowbuilding	2383	2,38	905	0,90	541	0,54	2	0,00	12	0,01	25	0,03	23	0,02	28	0,03
HighUrbanConstruction	147	0,15	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00
Rockstones, mines	0	0,00	217	0,22	24	0,02	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00
Together	52521	52,50	13786	13,78	6636	6,60	3713	3,71	3696	3,69	4408	4,41	5322	5,32	5333	5,33

Land use	Height, m. above sea level												Вместе			
	800-900		900-1000		1000-1100		1100-1200		1200-1300		1300-1400					
	ha	%	ha	%	ha	%	ha	%	ha	%	ha	%				
1	2	3	4	5	6	7	8	9	10	11	12	13	16	17		
The Cities	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	1237	1,24
Pshchenitsa	367	0,37	105	0,11	1	0,00	0	0,00	0	0,00	0	0,00	0	0,00	49979	49,96
Gardens	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	1123	1,12
Vineyards	4	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	528	0,53
Pastures	1530	1,53	384	0,38	217	0,22	98	0,10	28	0,03	1	0,00	12394	12,39		
Green grounds	189	0,19	3	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	6511	6,51
Power	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	4708	4,71
DryDriedLoose Valleys	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	673	0,67
Coniferous forests	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	535	0,53
Deciduous forests	617	0,62	260	0,26	334	0,33	137	0,14	103	0,10	30	0,03	8451	8,45		
Mixed forests	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	216	0,22
Marshes	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	844	0,84
Reservoirs	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	27	0,03
Stavy	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	25	0,02
Piece of lake	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	80	0,08
Bays	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	1137	1,14
Shrubs	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	2668	2,67
The soil layer without vegetation	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	248	0,25
Rocks	21	0,02	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	3827	3,83
Rural low construction	180	0,18	22	0,02	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	527	0,53
Citylowbuilding	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	3919	3,92
HighUrbanConstruction	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	147	0,15
Rockstones, mines	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	0	0,00	241	0,24
Together	2909	2,91	775	0,77	551	0,55	236	0,24	131	0,13	31	0,03	100047	100,0		

Table8 . Landuse according to grades of the slope of the terrain

Land use	Classes slopes%										Together			
	0-6		6-10		10-18		18-27		27-37					
	ha	%	ha	%	ha	%	ha	%	ha	%				
The Cities	49085	49,06	533,2	0,53	348	0,35	13	0,01	1	0,00	49979	49,96		
Pshchenitsa	1113	1,11	105,3	0,11	19	0,02	0	0,00	0	0,00	1237	1,24		
Gardens	1083	1,08	32,8	0,03	8	0,01	0	0,00	0	0,00	1123	1,12		
Vineyards	371	0,37	51,2	0,05	90	0,09	16	0,02	0	0,00	528	0,53		
Pastures	2895	2,89	2602,5	2,60	5956	5,95	932	0,93	8	0,01	12394	12,39		
Green grounds	658	0,66	1462,2	1,46	3486	3,48	893	0,89	12	0,01	6511	6,51		
Power	4632	4,63	53,8	0,05	19	0,02	3	0,00	0	0,00	4708	4,71		
DryDriedLoose Valleys	642	0,64	5,8	0,01	12	0,01	12	0,01	0	0,00	673	0,67		
Coniferous forests	535	0,53	0,0	0,00	0	0,00	0	0,00	0	0,00	535	0,53		

Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts).

Deciduous forests	595	0,60	1565,0	1,56	4655	4,65	1501	1,50	134	0,13	8451	8,45
Mixed forests	208	0,21	1,7	0,00	6	0,01	11	0,00	0	0,00	216	0,22
Marshes	844	0,84	0,0	0,00	0	0,00	0	0,00	0	0,00	844	0,84
Reservoirs	27	0,03	0,0	0,00	0	0,00	0	0,00	0	0,00	27	0,03
Stavy	25	0,02	0,4	0,00	0	0,00	0	0,00	0	0,00	25	0,02
Piece of lake	78	0,08	1,9	0,00	0	0,00	0	0,00	0	0,00	80	0,08
Bays	735	0,74	146,5	0,15	179	0,18	74	0,07	2	0,00	1137	1,14
Shrubs	2550	2,55	25,3	0,03	62	0,06	31	0,03	0	0,00	2668	2,67
The soil layer without vegetation	196	0,20	16,8	0,02	24	0,02	11	0,01	0	0,00	248	0,25
Rocks	310	0,31	585,5	0,59	2051	2,05	872	0,87	8	0,01	3827	3,83
Rural low construction	354	0,35	117,1	0,12	55	0,05	1	0,00	0	0,00	527	0,53
Citylowbuilding	3723	3,72	87,9	0,09	98	0,10	1	,01	0	0,00	3919	3,92
HighUrbanConstruction	147	0,15	0,0	0,00	0	0,00	0	0,00	0	0,00	147	0,15
Rockstones, mines	217	0,22	8,1	0,01	16	0,02	0	0,00	0	0,00	241	0,24
Together	71026	70,99	7403,2	7,40	17084	17,08	4369	4,37	165	0,17	100047	100,00

LITERATURE

- [1] Alekperov KA, 1980. Soil-erosion map for the protection of lands. Moscow.
- [2] Aliev G.A. the problem of desertification in Azerbaijan, ways of solving it, Baku "Zia-Nurlan".
- [3] Aliev GA, 1959. Soil of Azerbaijan. Publishing house Volobuev V.R. «AN. Azerb. The USSR "p. 122.
- [4] Aliev BG, Aliev ZG and Aliev IN, 2000. Problems of erosion in Azerbaijan and ways to solve it. Baku: ZIMA-CPI "Nurlan" p. 122.
- [5] Armand, DL, 1956. Anthropogenic erosion processes. In the book. "Agricultural erosion and the fight against it", ed. An USSR, Moscow.
- [6] Babaeva KM, 1995. Influence of simple and complex mineral fertilizers and lucerne sowing on restoration of fertility of eroded soils of the south-eastern slope of the Greater Caucasus. Author. Dis. Baku.
- [7] Bennet H.H., 1958. Fundamentals of soil protection (translated from English).
- [8] Grossgeim AA, 1948. Vegetative cover of the Caucasus. Publishing house of the Moscow Society of Naturalists, Moscow.
- [9] Huseynov AA, 1991. Efficiency of surface improvement of eroded pastures. All-Union Scientific Conference. Dushanbe, p.
- [10] MN Zaslavsky, 1969. To the draft classification of soils according to the degrees of erosion. Sat. "Materials on the methodology of soil-erosion mapping of erosion control measures." Moscow.
- [11] M. Zaslavsky. 1983. Erosiology. "high school".
- [12] Ibragimov, AA, Mapping of eroded soils on agricultural lands, for example, Dashkesan district, Azerb. THE USSR.
- [13] Ibragimov AA, 2000. Agroecological feature of eroded soils in Azerbaijan.
- [14] Materials on the study of erosion and irrigation and soil conservation in Azerbaijan. Baku.
- [15] Ibragimov A, A., 1972. Mapping of eroded soils on agricultural lands (on the example of the Dashkesan district of the Azerbjessian Soviet Socialist Republic). Questions of the methodology of soil-erosion mapping
- [16] Groom G, Mücher CA, Margareta I, Thomas W (2006) Remote sensing inlandcape ecology: experience and perspectives in the European context. Landscape Ecology 21 (3): 391-408.
- [17] Harris A, Carr AS, Dash J (2014) Remote sensing of vegetative integumentary dynamics and stability in southern Africa. International Journal of Earth Observation and Geoinformation 28: 131-139.
- [18] Klemas V (2013) The use of remote sensing for the selection and monitoring of wetland restoration sites: an overview. Journal of Coastal Research 29 (4): 958-970.

Citation: Dr.Z.H.Aliyev (2018). “ Spatial Analysis of Data on the Basis of the Digital Model of Relief and Locality (Example of Shemakha, Akhsu and Ismayilli Districts). ”. International Journal of Research in Agriculture and Forestry, 5(8), pp.29-35

Copyright: © 2018 Dr.Z.H.Aliyev et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.