

Effect of Supplemental *Plumeria Rubra* Leaf Meal on Performance of Starter Broiler Birds

Patricio De Los Ríos^{1,2*}, Olueze, C.C³ and Ahaotu, E.O³

¹Universidad Católica de Temuco, Facultad de Recursos Naturales, Departamento de Ciencias Biológicas y Químicas, Casilla 15-D, Temuco, Chile

²Núcleo de Estudios Ambientales, UC Temuco

³Department of Animal Production Technology, Imo State Polytechnic Umuagwo, Nigeria

***Corresponding Author:** Patricio De Los Ríos, Universidad Católica de Temuco, Facultad de Recursos Naturales, Departamento de Ciencias Biológicas y Químicas, Casilla 15-D, Temuco, Chile, Email: oshotex2003@yahoo.com.hk

ABSTRACT

Ninety "Ross" unsexed day old broiler chicks were used to determine the optimum level of *Plumeria rubra* leaf meal supplementation in broiler diet. The experiment was laid out in a completely randomized design (CRD). The birds were randomly selected and divided into five treatment groups of eighteen birds each. Each treatment group had two replicates with nine birds per pen. The birds were fed five diets containing *Plumeria rubra* leaf meal at 0, 0.5, 1, 1.5 and 2 % levels of inclusion. Results revealed that feed intake of birds on diet 1 (129.39g), 2 (131.71g) and 3 (128.24g) were similar but were significantly ($P < 0.05$) higher than those of birds fed diet 4 (126.56g) and 5 (110.60g). Similarly weight gains of birds on diet 1 (40.82g), 3 (40.20g) and 4 (39.80g) did not differ significantly ($P > 0.05$) but birds on diet 2 (44.80g) had a higher weight gain which resulted in a significantly higher final body weight. Haemoglobin (Hb) values of birds fed diet 2 (10.8g/dl) and 5 (10.2g/dl) were similar ($P > 0.05$) but significantly ($P < 0.05$) higher than those of birds fed diet 1 (8.9g/dl), 3 (8.8g/dl) and 4 (9.0g/dl). Also the packed cell volume (PCV) values of birds fed diet 2 (39%) was significantly ($P < 0.05$) higher than those fed diet 1 (32%), 3 (30%), 4 (34%) and 5 (36%) respectively. The overall result shows that birds on diet 2 (0.5% PRLM) had significantly ($P < 0.05$) superior body weight gain and final body weight than birds on diet 1 (0% PRLM), 3 (1% PRLM), 4 (1.5% PRLM) and 5 (2% PRLM). Also haematological values investigated were within accepted normal range for poultry, except for haemoglobin where diet 1 and 3 fell slightly below the recommended values for broiler birds.

Keywords: *Plumeria rubra*, Leaf meal, Performance, Starter Broiler Birds

INTRODUCTION

Nutrition is the most important consideration in any livestock enterprise. Its (livestock industry) survival is dependent on the availability of feedstuffs, which are mainly components of human food (Adewale *et al.*, 2018). The aim of keeping livestock and poultry is for the production of high quality protein. Agriculturists and Nutritionist in Nigeria have generally agreed that poultry is the fastest means of bridging the protein deficiency gap prevailing in the country (Ogba *et al.*, 2020 and Ahaotu, 2018).

Broiler birds are fast growing birds and are described as good converters of feed and are marketed from eight to twelve weeks. Broiler birds are also regarded as the type of birds that have high feed consumption and conversion ratio of non-conventional feed ingredients that

cannot be directly consumed by man into high quality meat which are needed in large quantity by man (Is-Haaq *et al.*, 2018). However, the production of broiler birds as meat bird cannot be effective in the absence of adequate feed and feed ingredients in right proportion. The productivity of poultry in the tropics has been limited by scarcity and consequent high prices of the conventional protein and energy sources. Protein sources are especially limiting factors in poultry feed production in the tropics (Ahaotu and Akagha, 2020, Ihezuo *et al.*, 2013).

It has been further estimated that 70-80% of poultry production inputs is been attributed to feed cost, hence making it imperative that good quality feed be available at reasonable price to make poultry production more profitable (Ebochuo *et al.*, 2017). Thus, the birds can only perform economically well and profitably if it

consumes, on daily basis, the appropriate amount of energy, protein, vitamins and minerals (Alionye *et al.*, 2020).

Antibiotics are drugs such as penicillin, amoxicillin and tetracycline that are used to kill or inhibit the growth of harmful bacteria (Ahaotu *et al.*, 2018b). Thus, the use of antibiotics in improving the rations of birds has been carried out by various authors. Ahaotu *et al.*, (2012b) reported that growth depression in broiler chickens fed high fibre diet was effectively and economically alleviated by the action of the following antibiotics namely tylosin, streptomycin and neo-terramycin. Also Ayo-Enwerem *et al.*, (2013) reported that the growth performance of broiler chicks fed a high fibre diet supplemented with antibiotics was better than those fed un-supplemented diet. Thus, antibiotics supplementation is an effective and readily adaptable strategy, if not for the possibility of their immunosuppressive actions under prolonged usage.

However the inclusion of antibiotics in livestock rations as feed additives is being discouraged because of the negative effect on the animals and the humans who consume the final animal products (Ogbuokiri *et al.*, 2015). This was also confirmed by Ahaotu *et al.*, (2018d) who reported the problems of residual effect in livestock products. Thus, there is the need to evaluate potential antibiotic alternatives to improve disease resistance in intensive food animal production. Nutritional approach to counteract the debilitating effects of stress and infection may provide products with useful alternatives to antibiotics. Improving the disease resistance of animals grown without antibiotics can benefit the animal's health, potentially increasing production efficiency and food safety. Improving disease resistance in food animals particularly in the absence of antibiotics treatment is a key strategy in the effort to increase food safety (Ezeafulukwe *et al.*, 2017). Hence, there is a need to search for locally available alternative sources of protein for use as feed supplement to poultry. One possible source of cheap protein to poultry is the leaf meal of some tropical plants (Gopi *et al.*, 2011).

Leaf meals of various plants have been incorporated in the diets of poultry as a means of reducing the high cost of conventional protein sources (Ayo –Enwerem *et al.*, 2018). Ahaotu *et al.* (2011) stated that leaf meals do not only serve as protein source but also provide some necessary vitamins, minerals and oxycarotenoids

which cause yellow colour of broiler skin, shank and egg yolk.

Among various types of *Plumeria* species, *Plumeria rubra* is the most economically important specie (Ihezuo *et al.*, 2013). It is widely distributed in Nigeria (Ahaotu *et al.*, 2018a) and is used as vegetable food for human consumption and animal feed resources during dry period (Ahaotu, 1997). The edible parts of the *Plumeria rubra* trees are exceptionally nutritious (Ahaotu *et al.*, 2018a). Leaf parts are promising as a food source in the tropics because the tree is full of leaves during the dry season when other foods are typically scarce (Ucan *et al.*, 2001). Recent studies conducted by Ogunwande *et al.* (2015) indicated that leaves of *Plumeria rubra* are rich in protein (28.2-36.2%) and contain considerable amounts of essential amino acids. The leafy part of *Plumeria rubra* could thus be used as a protein supplement for poultry.

This study therefore determined the haematology and blood chemistry of starter broilers fed graded levels of *Plumeria rubra* leaf meal and investigated the effects of feeding different levels of *Plumeria rubra* leaf meal (PRLM) on nutrient intake and growth performance of starter broilers.

At a high inclusion level, leaf meals lead to depressed growth. This may be attributed to their low digestibility and inadequate metabolisable energy content. Also with older broiler chicks, Is-Haaq *et al.* (2018) showed that diets containing 100g leaf meal per kilogram diet significantly reduced growth without affecting dry matter intake.

MATERIALS AND METHODS

Experimental Site

The experiment was carried out at the Poultry unit of the Teaching and Research farm, Imo State Polytechnic Umuagwo, Imo State. Umuagwo is in the derived savannah zone of Nigeria. The study area is located on latitudes 5° 28' 00" N and 5° 30' 00" N and longitudes 7° 01' 00" E and 7° 03' 00" E. The mean annual rainfall is 1247mm with a relative humidity of between 75 and 95%. It is situated at about 600 m above sea level with a mean annual temperature of 26.2°C (IMLS, 2009).

Preparation of Experimental Diets

Plumeria rubra leaves were collected from the Poultry unit of Imo State Polytechnic Umuagwo. Branches from matured *Plumeria rubra* trees, were cut, twigs and leaflets were removed by hand.

Effect of Supplemental *Plumeria Rubra* Leaf Meal on Performance of Starter Broiler Birds

The collected leaves were then spread on a (thinly sliced dried bamboo stalks and woven together like wire mesh) and dried in the open air until the leaves dried out uniformly. Using thinly sliced dried bamboo stalks and woven together like wire mesh for drying purpose was advantageous because it efficiently enhances air circulation during the drying process and prevents uneven drying of leaves by retaining the greenish color of the leaves. The dried

leaves were hand-crushed and further ground using mortar and pestle at 2 mm size.

Experimental Diets

Five experimental diets were used in the study. Treatment 1 (Diet 1) served as the control with 0% of *Plumeria rubra* leaf meal, while diets 2, 3, 4 and 5 contained *Plumeria rubra* leaf meal at 0.5, 1, 1.5 and 2% level of supplementation respectively (Table 1).

Table 1. Percentage Composition of the Experimental Diet

Ingredients	Diets				
	1	2	3	4	5
<i>Plumeria rubra</i> leaf meal	0	0.5	1	1.5	2
Maize	44.50	44.40	44.20	44.10	44.10
Wheat offal	13.80	13.70	13.60	13.40	13.10
Soya bean meal	9.40	9.30	9.20	9.20	9.10
Groundnut cake	14.20	14.10	14.10	14.00	14.00
Palm kernel cake	7.50	7.40	7.30	7.20	7.10
Fish meal	5.70	5.70	5.70	5.70	5.70
Blood meal	1.90	1.90	1.90	1.90	1.90
Bone meal	2.00	2.00	2.00	2.00	2.00
Salt	0.25	0.25	0.25	0.25	0.25
Vitamin premix	0.25	0.25	0.25	0.25	0.25
Lysine	0.25	0.25	0.25	0.25	0.25
Methionine	0.25	0.25	0.25	0.25	0.25
Total	100	100	100	100	100
Calculated Analysis					
Crude protein (%)	23.31	23.30	23.32	23.33	23.34
Energy (Kcal/k)	2,785.69	2,788.42	2,762.79	2,752.39	2,743.72
Crude fibre (%)	4.25	4.29	4.31	4.34	4.36
Determined Analysis					
Crude protein	24.06	24.28	23.62	24.53	24.65
Crude fibre	4.00	5.00	5.50	5.80	5.90
Ash	10	12	10	10	10
Ether extract	6	8	8	10	12
Moisture	10	10	12	10	12
NFE	45.94	40.72	40.88	39.67	37.45

Management of Birds

Few days to the arrival of the chicks, the brooding pen, the drinkers and the feeders were cleaned and disinfected. Wood shavings were spread on the floor. The hovers, feeders and drinkers were placed in position. Kerosene stoves and lanterns were used to provide the heat needed to keep the temperature within optimum range (between 33°C to 35°C). Feed and water were provided ad-libitum. Feeding was carried out twice, in the morning between the hours of 7 - 8am and in the evening between the hours of 5-6pm daily.

Response Parameters

Average Daily Feed Intake per Bird

This was calculated as the amount of feed left over subtracted from the total amount of feed

supplied. The result was then divided by the number of days of the experiment (28 days).

Average Daily Weight Gain Per Bird

This was estimated as the total weight of the birds at the end of the period (four week) less the total weight of the birds at the beginning of the study, divided by the number of days (28 days) and the result was then divided by the number of birds per replicate.

Feed Conversion Ratio

This was calculated as the average daily feed intake divided by the average weight gain.

Body Weight

This was measured on a weekly basis using a weighing balance. The birds were however weighed at the beginning of the experiment to determine their initial weight.

Haematological Studies

This was carried out using blood samples collected from each treatment. Two birds were collected from each replicate (four birds per treatment) for analysis. The parameters analyzed included; hemoglobin (Hb), packed cell volume (PCV), total bilirubin and aspartate amino-transferase (AST).

Statistical Analysis

At the end of the study (fourth week) data obtained from the response variables were subjected to analysis of variance using a one way (ANOVA) by Steel and Torrie (1980) in a completely randomized design (CRD).

Laboratory Analysis

Some quantity of the dried *Plumeria rubra* leaf meal which was obtained after processing was taken for analysis in the laboratory at Michael Opara University. Also some quantity of feeds formulated for the birds were analyzed in the laboratory that is diet 1, 2, 3, 4 and 5 containing 0, 0.5, 1.0, 1.5 and 2.0% levels of inclusion of *Plumeria rubra* leaf meal respectively.

Table2. Proximate Composition of *Plumeria rubra* leaf meal

Components	Percentage
Dry matter	89.70
Crude protein	24.06
Ash	6
Moisture	3.5
Crude fibre	12
Ether extracts	6

The result above showed that the *Plumeria rubra* leaf meal contains, 24.06% crude protein, 6% ash, 3.5% moisture, 12% crude fibre and 6% ether extract.

Table3. Composition of Essential Amino Acids of *Plumeria rubra* leaves (g/16g N)

Amino acids	<i>Plumeria rubra</i> leaves
Methionine	1.71
Cystine	2.04
Valine	5.34
Leucine	8.50
Phenylalanine	5.27
Threonine	4.45
Tryptophan	1.54
Isoleucine	4.47
Lysine	5.68
Arginine	6.4

Table4. Effects of Varying Levels of Supplemental *Plumeria rubra* leaf meal on the Performance of Broiler Birds

Parameters	Dietary Treatment					SEM
	(0%)	2(0.5%)	3(1%)	4(1.5%)	5(2%)	
Initial Body Weight (g).	70	73	75	72	74	0.61
Body Weight (g)	620 ^b	580 ^a	570 ^c	530 ^c	490 ^d	3.25
Avg. Daily Weight. Gain (g).	40.82 ^b	44.80 ^a	40.20 ^b	39.80 ^b	32.36 ^c	0.43
Avg. Daily Feed Intake (g)	129.39 ^a	131.71 ^a	128.24 ^a	126.56 ^b	110.60 ^c	1.74
Feed Conversion Ratio (g)	3.17 ^b	2.94 ^c	3.19 ^b	3.18 ^b	3.43 ^a	0.03

Average Daily Weight Gain (G)

Results presented in Table 4 shows that the highest daily weight gain (g) was observed in birds fed diet 2 (0.5% PRLM), while the lowest was observed in birds fed diet 5 (2% PRLM). It was observed that the body weight gain of birds decreases with increase in dietary levels of the PRLM. The result agrees with the observation made by Is-Haaq *et al.*, (2018), where graded dietary inclusion of *Pueraria phaseoloides* leaf meal induced cause related depression in growth of chicken even when maize oil was used to compensate for low metabolizable energy value of the leaf meal. Similar observation of decreasing performance of broilers at high levels of leaf meal inclusion have been observed by Mueller *et al.*, (2012) who reported that the body weight gain of broilers drastically reduced at high levels of *Plumeria rubra* leaf meal incorporation.

Also the work of Ahaotu *et al.* (2018e) follows the same suit with the incorporation of high levels of *Carica papaya* leaf meal. The effect observed above could have resulted from imbalance nutrient value and improper metabolism associated with the *Plumeria rubra* leaf meal as reported by Ihezuo *et al.* (2013). Other workers (Ahaotu *et al.*, 2018g) have also reported the effect of nutrient imbalance on monogastric animals fed high levels of unconventional feed ingredients. This could also be probably due to the effects of incomplete elimination of toxic factors as reported by Okonkwo and Ahaotu, (2014).

Average Daily Feed Intake

The analysis of variance conducted on the daily feed intake shows that there was significant (p<0.05) differences among the treatment means. The feed consumption of birds fed diet 1, 2 and 3 were similar (p>0.05) but significantly

($p < 0.05$) higher than those of birds fed diets 4 and 5. However the feed intake of birds fed diet 5 was significantly ($p < 0.05$) lower than that of birds fed diet 4.

The observed general trend was that of decreasing average feed intake as level of supplementation increased. This might be attributed to the bitter nature of the *Plumeria rubra* leaf which reduced the palatability of the feed. This is in consonance with the observations of Gupta *et al.*, (2007) who fed rabbits with raw *Plumeria rubra* seed cakes and reported reduced feed consumption among the rabbits on the test diets. Ahaotu *et al.* (2019) has also established the presence of bitter triterpenoids in the *Plumeria rubra* seeds. However, debitterization through water washing, alkali soaking and urea ammoniation to improve palatability has been recommended (Hossain and Becker, 2001). Also the decrease in feed intake observed could be as a result of anti-nutritional factors present in the test ingredient. This agrees with the findings of Kumar *et al.*, (2009) that incorporated different leaf meals noted for their anti-nutritional factors

in the diets of poultry, and observed decreases in feed intake.

Feed Conversion Ratio

The analysis of variance shows that there were significant ($P < 0.05$) differences among the treatment means. As shown in table 5, the amounts of feed consumed by birds fed diet 1 (0% PRLM), diet 3 (1% PRLM) and diet 4 (1.5 % PRLM) to gain the same unit (kg) of bodyweight, were similar ($P > 0.05$). However, birds fed diet 2 (0.5%, PRLM) were significantly ($P < 0.05$) higher than those birds fed diet 5 (2%) to gain the same unit of body weight. The efficiency of feed conversion of birds fed diets containing more than 0.5% PRLM were significantly ($P < 0.05$) lower when compared to those of birds fed diet 2 (0.5%, PRLM). The poor utilization of diets containing higher levels of PRLM might be related to the inability of the birds' enzyme to break down the active bitter substance in the PRLM and also the imbalance nutrient value and improper metabolism associated with the *Plumeria rubra* leaf meal as reported by Gupta *et al.* (2007).

Table 5. Effect of Supplemental *Plumeria rubra* Leaf Meal on the Hematological Values of Broiler Birds

Dietary Treatments Parameters	1	2	3	4	5	SEM
Hb (g/dl)	8.9 ^b	10.8 ^a	8.8 ^b	9.0 ^b	10.2 ^a	0.60
PCV (%)	32 ^c	39 ^a	30 ^c	34 ^b	36 ^b	1.85
Total Bilirubin (mg/dl)	2.35 ^a	2.15 ^{ab}	2.25 ^a	2.15 ^{ab}	2.05 ^{ab}	0.21
AST (Unit/l)	10.05 ^{ab}	10.15 ^a	10.25 ^a	9.98 ^{ab}	10.10 ^a	0.24

Haemoglobin (Hb) Values

The result of the hematological values of broiler birds revealed that there were significant difference ($P < 0.05$) among the treatments. Hb values of birds fed diets 2 and 5 were similar ($P > 0.05$) but significantly ($P < 0.05$) higher than those of birds fed diets 1, 3 and 4, which were themselves similar ($P > 0.05$). The results of the haemoglobin values of the birds on diet 3 and 1 fall slightly below the recommended range of 9-13g/dl as reported by Nkwocha *et al.*, (2018).

However, the values obtained for birds on diet 2, 4 and 5 fall within the accepted range as (7.2 - 11.4) reported by Okonkwo and Ahaotu (2014). This result implies that including PRLM in the diets of broilers as supplemental feed had little or no effect on the relative quantity of the complex iron containing the conjugate protein. This gives a clear-cut reason why the redness of the broiler bird's blood was not impaired.

Packed Cell Volume (PCV) Values

The result of the Packed Cell Volume (PCV) revealed that birds on diet 2, 4 and 5 were similar ($P > 0.05$) but significantly ($P < 0.05$) higher than those of birds fed diet 1 and 3, which were themselves similar ($P > 0.05$). The highest value of packed cell volume was obtained in birds on diet 2 (0.5% PRLM). Thus the values obtained for PCV across the treatment groups were within the normal range of 30- 40% as reported by Adeyeye *et al.*, (2013). This result implies that including PRLM in the diets of broiler birds as supplemental feed ingredient had little or no effect on the relative quantity of blood cells as compared with the total volume of blood (Onyekwere *et al.*, 2016).

Aspartate Aminotransferase (Ast) Values

The result of the hematological values for Aspartate amino transferase (AST) shows that, there was no significant difference ($P > 0.05$) among the treatments. The result obtained for AST across the treatment groups were within

the accepted normal range of 3-45 units/l as reported by Ayo – Enwerem *et al.*, (2017). This is also in accordance to the findings of Gupta *et al.*, (2007) who reported that usually low levels of AST are normally found in the blood of Poultry birds, but when high levels are found (400-4,000unit/1) then there is likely to be cases such as viral hepatitis and carbon tetrachloride poisoning.

Bilirubin Values

The result of the hematological values for total bilirubin shows that there was no significant difference ($P>0.05$) among the treatments. The low level of bilirubin obtained across the treatment groups is an indication of the normalcy of the blood level of the birds. High levels of bilirubin usually reveal that too much is being produced (usually due to increased destruction of red blood cells or that the liver is incapable of adequately removing bilirubin in a timely manner (due to blockage of bile ducts and liver diseases such as cirrhosis and acute hepatitis) (Ahaotu, 2011).

CONCLUSION

An experiment was conducted with Ninety “ROSS” unsexed day old chicks to determine the optimum level of supplementation of *Plumeria rubra* leaf meal in broiler diets. The experiment was laid-out in a completely randomized design (CRD). The birds were randomly allocated to five treatment groups of eighteen birds per treatment. Each treatment was replicated twice with nine birds per replicate. The five groups were randomly allocated to five diets containing varying levels of *Plumeria rubra* leaf meal (PRLM) of 0% (diet 1), 0.5% (diet 2), 1.0% (diet 3), 1.5% (diet 4) and 2.0% (diet 5). The birds' performances were measured and calculated on daily basis in terms of body weight gain, feed intake and feed conversion ratio. At the end of the experiment the haematological values were determined. The parameters determined for include hemoglobin (Hgb), aspartate aminotransferase (AST), bilirubin and packed cell volume (PCV).

The results of this study showed that there were significant ($P<0.05$) difference among the performance of birds across the various treatments in terms of average body weight gain, feed intake, and feed conversion ratio. Further supplementation of PRLM beyond 0.5% showed decrease in average bodyweight gain and average final weight of the birds. This study also showed that the supplementation of PRLM

had little or no effect on the hematological values of the birds, as the parameters evaluated for across the treatment groups fell within the accepted normal ranges for poultry birds.

The results of this study showed that the *Plumeria rubra* leaf meal supplementation in broiler diets could be effectively utilized at 0.5% to obtain reasonably good growth rate and reduced disease infection risk in the finished broiler birds.

REFERENCES

- [1] Adewale, I.T, Ahaotu, E.O, Onyekwere, M.U, Olaoye, J.C and Ikpe, J.N (2018). Serum Biochemical Indices and Hematological Profiles of broiler Birds Fed Varying Levels of Biscuit Waste Meal Based Diets. *Int J Anim Sci.* 2 (5): 1032-1035.
- [2] Adeyeye, S.A., Agbede J.O, Aletor V.A, Oloruntola O.D and E. O. Ahaotu (2017). Performance Of Weaner Rabbits Fed Diets Containing Graded Levels Of Processed Kolanut (*Cola Acuminata*) Pod Husks. *Proc. 6th ASAN-NIAS Joint Annual Meeting.* Sept 10-14, 2017. Pp 558-561.
- [3] Adeyeye, S.A, O.P.A. Oluwu, O. A. Adu, K. Sule, M. H. Ogunsipe, E. O. Ahaotu and J.O. Agbede (2013). Haematological and Biochemical studies of broiler chickens fed copper chloride supplemented diets. *Proc. 38th Conf., Nig. Soc. Anim. Prod.* 17-20 March, 2013. Pp 144-147.
- [4] Ahaotu EO and Akagha NU (2020). Factors Affecting Adoption of Indigenous Strategies for Climate Change Adaptation in Imo State, Nigeria (A Survey Using Systematic Sampling of Households in Orlu Local Government Area). *Acta Scientific Agriculture* 4 (1): 49-55.
- [5] Ahaotu, E.O, Nwoye, E.O and Lawal, M (2019). Biochemical Constituents, Economics of Production and Performance of Starter Broilers Fed Diets Supplemented with Natural Feed Additives. *International Journal of Animal and Veterinary Sciences.* 06 : 01-05
- [6] Ahaotu, E.O (2018a). Sheanut *Vitellaria paradoxa* cake as ingredient in the diet of West African Dwarf Sheep *Capra ovis*. *Aceh Journal of Animal Science.* 3 (2): 47-54. DOI: 10.13170/ajas.3.2.11325.
- [7] Ahaotu, E.O (2018b). Optimum Supplementation Levels of Low-and-High Cyanide Cassava Peels on the Performance, Nutrient Digestibility and Serum Metabolites of Growing Rabbits. *International Journal of Animal and Veterinary Sciences.* 05: 15-18.
- [8] Ahaotu E.O, Uwalaka R, Edih M.C and Ihiaha P.O (2018a). Water Purification Using *Moringa oleifera* (Horse Radish) Leaves and Seeds. *Journal of Agricultural, Biological and Environmental Sciences.* 05: 13-18.

- [9] Ahaotu E.O, Uwalaka R.E, Edih M.C and Ihiaha P.O (2018b).Effect of Herbal Tablet Prepared from *Moringa oleifera* Leaves Extract. Journal of Agricultural, Biological and Environmental Sciences. 5: 06-12.
- [10] Ahaotu E.O, Uwalaka R.E, Edih M.C and Ihiaha P.O (2018c).Characterization and Extraction of Oil, Biogas and Biodiesel from *Moringa oleifera* Seeds. Journal of Agricultural, Biological and Environmental Sciences. 5: 01-05.
- [11] Ahaotu, E. O, Akinfemi, A. and Okorie, K. C (2018d).Economic Importance and Widespread of Ectoparasites Infestation in Indigenous Chickens (*Gallus gallus domesticus*).A Study from Selected Local Government Councils and States in Nigeria. Direct Res. J. Vet. Med. Anim. Sci. 3 (4): 21-27.
- [12] Ahaotu, E.O., P. O. Ihiaha, C. O. Osuagwu and E.U. Nwabueze (2018e).Replacement value of Feather Meal for Fishmeal on the Performance of Guinea cock.Proc.7th ASAN-NIAS Joint Annual Meeting, 9th- 13th September, 2018, Ilorin, Kwara State, Nigeria. pp. 268 – 271
- [13] Ahaotu, E.O, Onyegbula, T and Ahaotu, Esther O (2018f). Haematological Characteristics and Organoleptic Test of Feeding Different Levels of Pawpaw (*Carica papaya*) Leaf Meal on Finisher Broiler Birds. International Journal of Animal and Veterinary Sciences. 05: 10-14.
- [14] Ahaotu, E. O, Amajioyi, N and Okorie, K. C (2018g). Sustainability of Guinea Fowl (*Numidia Meleagris*) Layers Fed varying Levels of *Centrosema Pubescent* Leaf Meal. Direct Res. J. Vet. Med. Anim. Sci. 3 (4): 28-35.
- [15] Ahaotu, E.O, Emeribe, E.O and Akinfemi, A. (2018h). Carcass and Performance Characteristics of Finisher Broiler Birds Fed Jack Bean (*Canavalia ensiformis*) Sievate Fortified with Exogenous Enzyme. Journal of Meat Science and Technology. 06 (2): 19-24.
- [16] Ahaotu, E.O and U.T Agunanne (2017).Egg Production Pattern In Anthony Patience Farms, Atta-Ikeduru, Imo State, Nigeria: Influences On The Phase Of Production And The Estimated Period Of Oviposition. Proc. 6th ASAN-NIAS Joint Annual Meeting. Sept 10-14, 2017. Pp 375-380.
- [17] Ahaotu, E.O, Ihiaha, P.O, Edih, M.C, Ayo-Enwerem, C.M and A. Akinfemi (2015). Effect of dietary substitution of Wheat Offal with Cattle Dung on Growth Performance and Organ Weight Characteristics of Starter Broilers. Proc. 28th Ann. Conf. Nig. Assoc. of Health and Techno.3-6 Nov, 2015. Pp 6-10.
- [18] Ahaotu, E.O., C. Vajrabukka, A.S. Adeyeye, A.U. Onuruka, V.I. Ehirim and F. N. Madubuike (2012a). Performance and carcass characteristics of starter broiler chicks fed high fiber rice milling waste diets. Animal Production Research Advances, 7(4): 286-289.
- [19] Ahaotu, E. O., M. C. Edih, A.U. Onuruka, V. I. Ehirim and A. Sirimongkolkasen (2012b). Effects of replacement of soybean (*Glycine max*) with pigeon pea (*Cajanus cajan*) in starter broiler ration. International Journal of Tropical Agricultural and Food Systems. 6(1): 40 – 44.
- [20] Ahaotu, E.O, U.D.E. Ogbuokiri, A.U. Korie; B.U. Ekenyem, C.I.F. Onwuka, I.C. Okoli, Peace .O. Njoku, E.C. Ndubuisi and F.N. Madubuike (2011). Effects of graded levels of pigeon pea meal on growth performance and Organ Characteristics of finisher broilers. Animal Production Research Advances. 7(2): 125 – 129.
- [21] Ahaotu, E.O. (2011). Effects of dietary substitution of rubber seed cake for groundnut cake on the gross morphology and body conformations of broiler birds. Animal Production Research Advances. 7(1): 69 – 73.
- [22] Ahaotu, E.O., K. Karsten, Peace .O .Njoku, N. Yang, B.U. Ekenyem, A.U. Korie and F.N. Madubuike (2009). Effects of partial replacement of Soya bean meal with cassava meal in broiler finisher rations. Animal Production Advances, 5(4): 295 – 299.
- [23] Akinfemi, A, Ososanya, T. O and Ahaotu, E.O (2018).Nutrient and Anti-Nutrient Components of Some Tropical Tree Seeds. Nigerian J. Anim. Sci., 20 (1): 115-122.
- [24] Alionye, E.B, Ahaotu, E.O, Ihenacho, R.O and Chukwu, A.O (2020). Comparative Evaluation of Swine Production With Other Domestic Livestock in Mbaitolu Local Government Area of Imo State, Nigeria. Sustainability, Agri, Food and Environmental Research. 8(4): 222-244. <http://dx.doi.org/10.7770/safer-V0N0-art1989>
- [25] Ayo –Enwerem, M.C, Ahaotu, E. O and Oji, U. 1 (2018).Potential Degradability and Chemical Composition of Six Selected Multipurpose Fodder Trees and shrubs of the humid zone. *Journal of Agricultural Sciences and Research*.1 (1): 43-50.
- [26] Ayo- Enwerem, C.M., C.F. Ezeafulukwe, E.O. Ahaotu and A. Akinfemi (2013).Effects of Replacement of Fish Meal with Spent Grain on the Performance of Growing Pigs. Inter J Vet Sci. 2 (3): 78 – 80.
- [27] Ebochuo V.C., Ebigbi K.G., Ahaotu E.O. and Singh V (2017). Growth Performance of Juvenile African Catfish Fed *Ageratum Conyzoides* Leaf Meal Based Diets. Journal of Agricultural, Biological and Environ Sciences. 04: 47-50.
- [28] Ezeafulukwe, C .F, L.C. Onuoha and E.O. Ahaotu (2017). Growth Performance of *Clarias gariepinus* (Buchell 1822) Fed Varying Inclusion Levels of Cassava Flour. Greener Journal of Biological Sciences Vol. 7 (5): 45-49.
- [29] GopiJ, Khatri P, Singh N, Gaud H and Patel R. (2011). Phytochemical and pharmacological potential of *Plumeria rubra* Linn. (Apocynaceae):

- A review. *International Journal of Pharmaceutical Science*. 3: 1162-1168
- [30] Gupta M, Mazumdar U.K and Gomathi P. (2007). Evaluation of antipyretic and antinociceptive activities of *Plumeria acuminata* leaves. *J Med Sci*. 7: 835-839.
- [31] Hossain, M.A and Becker, K. (2001). Nutritive value and anti-nutritive factors in different varieties of *Sesbania* seeds and their morphological fractions. *Food Chemistry*. 73: 421-431.
- [32] Ihezuo, J.P, Ahaotu, E.O and Uwalaka, R.E (2013). Replacement of Rice and Maize Stalks with *Tephrosia spp* and *Moringa oleifera* cakes in the diet of Small Ruminants raised under humid tropics. *Inter J Appl Sci Engr* 1(2): 66-68.
- [33] Ihenacho, R.A, Ahaotu, E.O and Ayo – Enwerem, C.M (2016). Performance of Pullet Growers fed Feather Meal Supplemented with Fish Meal. Book of Proceedings of Academic Conference on Positioning Sub-Saharan Africa for Development in the New Development 22-23rd June, 2016, Cambridge Hall, University of Ghana, Legon Campus, Accra. Vol. 9 (2): 216 – 221.
- [34] Imo State Ministry of Lands and Survey (IMLS) (2009). Longitudes and Latitudes locations of Umuagwo, Ohaji / Egbema Local Government Area of Imo State, Nigeria.
- [35] Is-Haaq, A.T, Ahaotu E.O, Onumajuru C.G, Akinfemi. A and Mako A. A (2018). Replacement value of *Puerairia phaseoloides* leaf meal for soya bean meal on the Performance and carcass evaluation of starter broilers. *Journal of Research in Microbiology and Biotechnology*. 1 (1): 5-9. DOI: 10.5281/zenodo.1403686
- [36] Kumar A, Chanda I and Singh A. (2009). Extraction and evaluation of pharmacological activity of saponins extract of *Plumeria rubra* leaves. *Pharmacology online*. 1: 969-974
- [37] Mueller K., Blum N.M., Kluge H and Mueller A.S. (2012). Influence of broccoli extract and various essential oils on performance and expression of xenobiotic- and antioxidant enzymes in broiler chickens. *British Journal of Nutrition*, 108, 588–602.
- [38] Nkwocha, G.A., K.U. Anukam, E.O. Ahaotu and O.I. Prudent (2018). Performance and Hematological Profile of Broiler Chicks on Toasted Tiger Nut (*Cyperus esculentus*) Treated With Bakery Yeast. *Proc. 43rd Annual Conference of the Nigerian Society for Animal Production, March 18th – 22nd 2018, FUT Owerri*, Nigeria. Pp 484 – 487.
- [39] Ogba, O, Ahaotu, E.O, Ihenacho, R.O and Chukwu, A.O (2020). Challenges of Small Poultry Farms in Layer Production in Ikwuano Local Government Area of Abia State, Nigeria. *Sustainability, Agri, Food and Environmental Research*, 8(4): 212-222.
- [40] Ogbuokiri, U.D.E, M.C. Edih, V. Izuazu, E.O. Ahaotu and C.I. Okoli (2015). Effect of Infection with *E. Necatrix* on the Body Weight gains and Spermatogenesis in Broiler Breeder Males. *International Journal of Tropical Agriculture and Food Systems*. 9: 12-15.
- [41] Ogunwande I.A, Opoku A.R and Oladipupo A.L, (2015). Chemical composition of essential oils of *Plumeria rubra* L. grown in Nigeria. *European J Med Plants*. 6 (1): 55- 61.
- [42] Okonkwo, V.N and E.O. Ahaotu (2014). Effects of cassava pulp sievate meal on hematological indices of finisher broiler birds. *World Journal of Agricultural Sciences*. 2 (6): 142 – 145.
- [43] Ononiwu, C.C., Ahaotu, E.O, V.N. Okonkwo, C.M. Ayo-Enwerem and H. Botchway (2017). Effects of Bambara Nut Sievate Fortified Exogenous Enzyme on Performance and Carcass Characteristics of Finisher Broiler Birds. *Proceedings of the 20th Biennial Conference, Ghana Society of Animal Production (GSAP)*, 1-5 August, 2017. Sasakawa Centre, University of Cape Coast, Ghana Pp 73-80.
- [44] Onu, P.N., F.N. Madubuike, B.U. Ekenyem and E.O Ahaotu (2010). Effect of enzyme supplementation of heat-treated sheep manure diets on nutrient digestibility of finisher broilers. *Proceedings of the 15th Annual Conference of the Animal Science Association of Nigeria (ASAN)*, Sept. 14 – 16, 2010, Uyo, Nigeria.
- [45] Onu, P.N, M.O. Otuma, P.E. Nwakpu and E.O. Ahaotu (2008). Enzyme and Probiotic Supplementation of Maize processing waste based diets for weaned Rabbits. *Proc. 42nd Annual Conf. Agricultural Society of Nigeria (ASN)*. October 19th – 23rd 2008. Ebonyi State University, Abakiliki, Nigeria.
- [46] Onyek were, M.U, Okechukwu, S.O., Olabode, A.D and Ahaotu E.O (2017). Serum Testosterone and Hormonal Profile of Rabbits Fed Raw Garlic rhizome flour meal (*Allium sativum*). *Proc. 6th ASAN-NIAS Joint Annual Meeting*. Sept 10-14, 2017. Pp 167-170.
- [47] Onyekwere M.U., Okechukwu, S.O., Jiwuba, P. C., Onwujiariri, E.B. Ahaotu, E.O and Anukam K.U (2016). Effect of dried cocoyam tubers (*Colocasia esculenta*) meal on growth performance and blood serum mineral profile fed broiler starter. *Proc. 21st Ann. Conf. Animal Science Association of Nigeria*. Porthacourt, Rivers State. 18th -22nd Sept, 2016. Pp 184 - 189.

Effect of Supplemental *Plumeria Rubra* Leaf Meal on Performance of Starter Broiler Birds

- [48] Steel R. G. D and Torrie, J. H. (1980). *Principles of Statistics: A Biometric Approach*. 2nd Edition, McGraw Hill Book Company Incorporated, New York, USA.
- [49] Ucan, B. Cokun, V. Kurtolu and Cetingul, I.S (2001). Effect of dietary probiotic on performance and humoral immune response in layer hens. *Br. Poult. Sci.*, 42: 456-461

Citation: *Patricio De Los Ríos , Olueze, C.C and Ahaotu, E.O, “Effect of Supplemental Plumeria Rubra Leaf Meal on Performance of Starter Broiler Birds”, International Journal of Research in Agriculture and Forestry, 7(2), 2020, pp 15-23.*

Copyright: © 2020 Patricio De Los Ríos. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.